


# Distribution of Counties with Laurel Wilt Disease\* by year of Initial Detection

\* Laurel Wilt Disease is a destructive disease of redbay (*Persea borbonia*), and other species within the laurel family (*Lauraceae*) caused by a vascular wilt fungus (*Raffaelea lauricola*) that is vectored by the redbay ambrosia beetle (*Xyleborus glabratus*). The pathogen has been confirmed through laboratory analysis of host samples collected in the counties highlighted.




Initial Detection of *Xyleborus glabratus* - May 2002  
Port Wentworth, GA


Updated: January 30, 2015


- 2004
- 2005
- 2006
- 2007
- 2008
- 2009
- 2010
- 2011
- 2012
- 2013
- 2014
- 2015


### Information Provided By:


 **Chip Bates**  
cbates@gfc.state.ga.us


 **Laurie Reid**  
lreid@forestry.state.sc.us

 **Rob Trickel**  
rob.trickel@ncagr.gov

 **John J. Riggins**  
jriggins@entomology.msstate.edu

 **Jeff Eickwort**  
Jeffrey.Eickwort@freshfromflorida.com

 **Brent Cutrer**  
mcutrer@ldaf.state.la.us

 **Dana McReynolds Stones**  
Dana.McReynolds@forestry.alabama.gov

