

#### **Edible Flowers**

Fact Sheet 7.237

Gardening Series | Flowers

By S.E. Newman and A.S. O'Connor\* (10/20)

Edible flowers have been used in the culinary arts for flavor and garnish for hundreds of years. Early reports indicate that the Romans used flowers in cooking, as did the Chinese, Middle Eastern, and Indian cultures. During Queen Victoria's reign, edible flowers were popular. They have maintained their popularity today.

Many flowers are edible and the flowers of most culinary herbs are safe. However, proper identification of the plant is essential because some flowers are poisonous and should not be eaten (Table 1). Many plants have similar common names, which may cause added confusion. Always use the scientific name when choosing a flower.

Pick flowers early in the day. Use them at their peak for the best flavor. Avoid unopened blossoms (except daylilies) and wilted or faded flowers, which may have a bitter or unappealing flavor. Do not use flowers that have been sprayed with pesticides or collect flowers from plants that have been fertilized with raw manure. Generally avoid purchasing flowers from florists, garden centers, or nurseries since these flowers are not grown for consumption. Table 2 lists many plants that can be added to food for flavor, aroma, color, or garnish.

Fresh flowers also can be preserved for later use. Choose flowers with larger petals, like pansies, and paint the petals with an egg-white wash. Use a soft brush and pasteurized dehydrated egg whites to avoid food-borne illness. After painting, dust the petal with super-fine granulated sugar and dry it. Store preserved flowers

in an airtight container in a cool, dark place. Avoid dark-colored petals; they turn even darker with this treatment.

# **Using Edible Flowers**

To avoid stomach upset or to determine if there is an allergic reaction, limit consumption to small tastes. Slowly incorporate large quantities into the diet. Edible petals or entire flowers can be eaten. However, remove stems, anthers and pistils because they may be bitter (Figure 1). Use flowers that are free of insects and disease. Many edible flowers are high in vitamin C and/or vitamin A, along with other essential nutrients. Use them as garnishes and in salads. Recipes for flowers may be found in the following areas: baking, sauces, jelly, syrup, vinegars, honey, oil, tea, flowerscented sugars, candied flowers, wine, and flavored liquors. Flavored vinegars and oils prepared at home have a limited shelf-life and should be stored in the refrigerator (Kendall and Rausch, 2012). Pick fresh flowers and gently rinse with cool running water. Placed washed flowers between damp paper towels. Refrigerate until ready to use. Some flowers may have an extended shelf life if washed just before use. Flowers and herbs may also be dried. Refer to "Herbs: Preserving and Using" for more information.




Figure 1: Flower parts.



## **Quick Facts**

- Proper identification of edible flowers is important.
- Use flowers that are grown without pesticides.
- For best flavor, use flowers at their peak.
- Introduce new flowers into the diet slowly to be able to pinpoint allergic reactions or stomach upset.
- Edible flowers also may be preserved in oils or vinegars.

\*S.E. Newman, Colorado State University Extension greenhouse crops specialist and professor, horticulture and landscape architecture; and A. Stoven O'Connor, Extension horticulture agent, Larimer County. 9/13. Revised 10/20.

Table 1: Some common house and garden plants with toxic plant parts or toxic flowers (not a complete list, if in doubt, consult a poisonous plant reference).

Scientific name	Common names	Scientific name	Common names
Achillea millefolium	Yarrow, milfoil	Lantana camara	Lantana, red sage, shrub verbena
Anconitum spp.	Monkshood	Lathyrus spp.	Sweet pea (seeds)
Cestrum spp.	Day blooming jasmine, Night blooming jasmine		
Clematis spp.	Clematis, virgin's bower	Lobelia spp.	Cardinal flower
Colchicum spp.	Crocus	Narcissus spp.	Daffodil, jonquil
Convallaria majalis	Lily of the Valley	Nerium oleander	Oleander
Daphne mezereum	Daphne	Nicotiana spp.	Flowering tobacco
Datura spp.	Jimson weed	Papaver somniferum	Opium poppy, common poppy
Delphinium spp.	Larkspur	Phoradendron spp.	Mistletoe
Dicentra formosa	Bleeding heart	Physalis spp.	Chinese or Japanese lantern
Digitalis purpurea	Foxglove, digitalis	Rhododendron spp.	Azaleas, rhododendrons, rose bay
Euphorbia spp.1	Euphorbia	Ricinus communis	Castor bean, African coffee tree
Hippeastum spp.	Amaryllis	Zantedeschia aethiopica	Calla lily
Hyaninthus spp.	Hyacinth	Zigadenus sup.	Death camas, alkaligrass, wild onion

Poinsettias are not considered poisonous, but they are not edible. If eaten, all plant parts may cause varying degrees of mouth irritation, and vomiting, but not death. The cultivated rubber tree Heavea brasiliensis, Manioc or cassava (Manihot), and Castor bean (Rincinus) are close relatives, which are poisonous. With its close genetic ties to the rubber tree, which is where natural latex is derived, those who are sensitive may also be sensitive to the latex from poinsettias.

Table 2: Recommended plants with edible flowers.

Plant name	Type <sup>1</sup>	Flower color	Bloom	Taste	Fragrance	Comments and hints
Abelmoschus aesculentus Okra, gumbo, gombo	А	Yellow, red	Mid-July to August	Mild, sweet and slightly mucilaginous	_	Full sun, hot weather; prefers clay to clay loam.
Agastache foeniculum Anise hyssop	Р	Lavender	July to frost	Strong anise, sweet, licorice	-	May die back to the ground; readily resows; full sun to light shade; well-drained soil.
Alcea rosea Hollyhock	Р	Various	July to frost	Little, slightly bitter	-	Best as a garnish or container for dip. Full sun to light shade.
Allium schoenoprasum Chive	Р	Lavender, red to purple	May to June	Onion, strong	Onion	Separate florets to serve. Forms clumps; part shade to full sun; indoors.
Allium tuberosum Garlic chive	Р	White	August to frost	Onion, strong	Onion	Separate florets to serve. Partial shade to full sun; also indoors.
Anethum graveolens Dill	А	Yellow	June to frost	Stronger than leaves	Dill	Resows readily, tolerates poor soil but prefers well-drained soil; full sun.
Anthemis nobilis English chamomile	P	White petals; yellow center	Late June to frost	Sweet apple flavor	_	Ragweed sufferers may be allergic to chamomile; drink no more than one cup of tea per day. Prefers moist, well-drained soil; full sun to part shade.
Anthriscus cerefolium Chervil	А	White	May to June	Parsley-like, hint of citrus, tarragon	-	Start in cold frame.
Begonia x tuberhybrida Tuberous begonia	TP	White, pink, yellow, red, orange and combinations	July to August	Citrus	_	Grow indoors or out; dig tubers each fall, just after frost and store; prefers
Bellis perennis English daisy	Р	White to purple petals	April to September	Mild to bitter	_	Use as garnish or in salads. Thrives in cold weather; prefers full sun, moist soil.
Borago officinalis Borage	А	Blue, purple to lavender	June to July	Cucumber	-	Use as garnish; may be candied. Full sun; light, poor, dry soil. Attracts bees.
<i>Brassica</i> spp. Broccoli, cauliflower	B (grown as A)	Buds: blue- green; yellow flower, white buds in cauliflower	June to August	Broccoli	-	Vegetable. Prefers full sun; rich, well-drained soil. Sow indoors six weeks prior to transplant.
Brassica spp. Mustard	А	Yellow	April to May	Mustard, hot	_	Salad garnish. Watch for allergies.

#### (Continued)

Table 2: Recommended plants with edible flowers.

Plant name	Type <sup>1</sup>	Flower color	Bloom	Taste	Fragrance	Comments and hints
Calendula officinalis Calendula, pot marigold	A	Yellow, gold/ orange	June to August	Tangy and peppery	-	Ornamental. Dries well. Prefers cool weather; rich loam; direct sow.
Calendula officinalis Calendula, pot marigold	A	Yellow, gold/ orange	June to August	Tangy and peppery	_	Ornamental. Dries well. Prefers cool weather; rich loam; direct sow.
Carthamus tinctorius Safflower, American safflower, saffron	A	Yellow to deep red	August	Bitter flavor	_	May impart yellow color to cooked foods. Full sun; light, dry, well-drained soil; start indoors and transplant.
Centaurea cyanus	A	White, pink, blue	May to frost	Mild	_	Annual; petals are edible; the calyx is bitter
<i>Cercis canadensis</i> Redbud	Р	Pink	April	Beanlike to tart apple	_	Native tree to U.S.; may be marginal in Colorado. Full sun to part shade; sandy loam; difficult to transplant.
Chrysanthemum coronarium Garden chrysanthemum, shungiku	Р	Yellow to white	August to October	Mild	-	Ornamental. Full sun; rich, moist, well-drained soil.
Chrysanthemum leucanthemum Oxeye daisy	Р	White/yellow center	April to August	Mild	_	Use as garnish or in salad. Full sun; rich, moist, well-drained soil, invasive.
Cichorium intybus Chicory	Р	Blue to lavender	July to frost	Pleasant, mild-bitter similar to endive	-	Considered a weed. Grows in most soils; sun to shade; invasive.
<i>Citrus limon</i> Lemon	Р	White	Varies with variety	Citrus, slightly bitter	Sweet	Subtropical tree but may be grown indoors.
Citrus sinensis Orange	Р	White	Varies	Citrus, sweet/strong	Perfume, sweet	Subtropical tree but may be grown indoors.
Coriandrum sativum Coriander	А	White	June to frost	Like leaf but milder	Fragrant	Herb. Sow continuously for several harvests; sun; rich, well-drained soil.
<i>Cucurbita</i> spp. Squash or pumpkin	А	Orange, yellow	July to August	Mild, raw squash	Slightly floral	Vegetable. Enrich soil with compost; prefers full sun.
Cynara scolymus Artichoke	A	Immature head: green	Fall	Artichoke	=	Prefers rich soil, abundant moisture; propagate from division for annual harvest.
Dendranthema x grandiflorum Chrysanthemum	Р	Red, yellow, pink, orange, purple, white	August to October	Varieties differ, strong to bitter	Pungent	Ornamental. Full sun; rich, moist, well-drained soil.
<i>Dianthu</i> s spp. Dianthus or pinks	Р	Pink, white and red	June to August	Spicy, cloves	Some varieties are spicier	Ornamental. Tolerates wide range of soils; full sun.
<i>Eruca vesicaria</i> Rocket, arugula	А	White	May to frost	Nutty, smoky, less piquant than leaves	-	Salad green. Sow continuously for harvest; full sun to light shade; well-drained soil.
<i>Feijoa sellowiana</i> Pineapple guava	Р	White to deep pink	Grow indoors	Floral flavor; papaya or exotic melon	=	Grow indoors in a greenhouse. Rich, well-drained soil; full sun-light shade.
Foeniculum vulgare Fennel	Р	Pale yellow	July to August	Licorice, milder than leaves, sweet	-	Tolerates wide range of soils; part shade to full sun.
<i>Galium odoratum</i> Sweet woodruff	P	White	May	Sweet, grassy, vanilla	Vanilla	Herb or ground cover. May be invasive; prefers shade. Can have a blood thinning effect if eaten in large quantities. It is considered liver toxic. Best as a garnish or container for dips or spreads.
Gladiolus spp. Gladiolus	TP	Various except true blue	6-8 weeks after planting	Mediocre	_	Best as a garnish or container for dips or spreads.
Hemerocallis fulva Daylily	Р	Tawny orange	June to July	Cooked, combination of asparagus/zucchini	-	All parts are edible. Full to part shade; easy to grow. May Lilies contain alkaloids and are NOT edible. Day Lilies may act as a diuretic or laxative; eat in moderation.
Hibiscus rosa-sinensis Hibiscus, China rose, Rose-of-China	Р	Orange, red or purplish red	Late summer	Citrus/cranberry flavor	_	Subtropical tree but may be grown indoors. Used in many tea flavorings.

#### (Continued)

Table 2: Recommended plants with edible flowers.

Plant name	Type <sup>1</sup>	Flower color	Bloom	Taste	Fragrance	Comments and hints
Hibiscus syriacus Rose-of-Sharon	Р	Red, white, purple, violet	July to August	Mild, nutty	-	Shrub. Prefers well-drained soil; full sun to part shade; deadhead to keep blooming.
Hyssopus officinalis Hyssop	Р	Blue, pink, white	July to October	Bitter; similar to tonic	_	Used to flavor chartreuse, a liqueur; strong flavor. Prefers part shade and well-drained soil.
Lavandula angustifolia Lavender	Р	Lavender, purple pink, white	June to early August	Highly perfumed	Floral	Taste may be very strong depending on the plant.
Levisticum officinale Lovage	В	Yellow, white	August	Mild celery	-	Herb.
<i>Malu</i> s spp. Apple or crabapple	Р	White to pink	May	Slightly floral to sour	Sweet floral	Petals may be candied. Seeds are poisonous. Specimen tree, prefers full sun, fertile soil.
Melissa officinalis Lemon balm	Р	Creamy white	July to August	Lemony, sweet	Lemon	Herb. May be invasive.
<i>Mentha</i> spp. Mint	Р	Lavender, pink to white	July to September	Minty; milder than leaves	Fresh, minty	Herb. May be invasive; tolerates a wide range of soils; prefers part shade.
Monarda didyma Bergamot, bee balm, Oswego tea	Р	Red, pink, white, lavender	July to August	Tea-like, more aromatic than leaves	Sweet, perfumed	Attracts hummingbirds and butterflies; part shade to full sun; prefers moist, rich soil. Powdery mildew when grown in part shade.
Muscari atlanticum, M. botryoides, Grape hyacinth	Р	Pink, blue	April to May	Grapey, slightly sour with bitter aftertaste	Grapey	Bulb.
Ocimum basilicum Basil	А	White to pale pink	July to frost	Milder than leaves, spicy	-	Sow continuously for several harvests; well-drained rich soil; full sun.
<i>Origanum majorana</i> Marjoram	TP	Pale pink	June to August	Spicy, sweet	-	Herb. Prefers full sun and dry, alkaline, well-drained soil.
<i>Origanum</i> spp. Oregano	Р	White	June to August	Spicy, pungent-like leaves	-	Herb. Prefers full sun and dry, alkaline, well-drained soil.
Pelargonium spp. Scented geranium	TP	White, red, pink, purple	Varies	Like variety selected, e.g., rose, lemon; varies	Mild	Ornamental. Prefers full sun; light, dry well-drained soil.
Phaseolus coccineus Scarlet runner bean	TP	Bright orange to scarlet	July to August	Raw bean but milder	-	Vegetable. Flower crunchy; use in salad.
<i>Pisum sativum</i> Garden pea	А	White, tinged pink	May to June	Raw peas	_	Vegetable. Prefers full sun; sandy, well-drained soil.
<i>Poterium sanguisorba</i> Burnet	Р	Red	July to August	Cucumber	_	Salad herb. May be invasive; tolerates wide range of soil; sun or part shade.
<i>Prunu</i> s spp. Plum	Р	Pink to white	April to May	Mild, like flower	Sweet	Petals candy well. Pits of mature fruit are poisonous.
<i>Raphanus sativus</i> Radish	А	White, pink, yellow	One month after planting	Spicy	_	Prefers full sun; well-drained, sandy soil but will grow in almost any soil.
Rosa spp. Rose	Р	White, pink, yellow, red, orange	May to June, September	Highly perfumed; sweet to bitter	Rose	Ornamental. Remove sour petal base. Full sun; rich, well-drained soil.
Rosmarinus officinalis Rosemary	TP	Pale blue, dark blue, pink, whit	Depends on cultivar	Mild rosemary	Delicate	Herb. Do not cook flower. Tolerates full sun to part shade; well-drained, evenly moist soil.
Salvia elegans Pineapple sage	TP	Scarlet	September	Pineapple/sage overtones	-	Herb. Prefers full sun; light, well-drained soil; may be invasive.
Salvia officinalis Garden sage	Р	Blue, purple, white, pink	May to July	Flowery sage, slightly musky	-	Herb. Full sun to light shade; sandy, well-drained soil; may be invasive.
Satureja hortensis Summer savory	А	Pink	July to August	Mildly peppery, spicy	-	Herb. Prefers full sun; light, sandy soil.
Satureja montana Winter savory	Р	Pale blue to	July to August	Mildly peppery, spicy	_	Herb. Prefers full sun; light, sandy soil.
Syringa vulgaris Lilac	Р	White, pink, purple, lilac	April to May	Perfume, slightly	Lilac	Candies well. Prefers well-drained, alkaline soil; sun to part shade.
Tagetes erecta African marigold	А	White, gold, yellow, red	May to September	Variable; some cultivars are strong and bitter	Strong, pungent	Ornamental. Prefers full sun; well-drained soil.
Tagetes tenuifolia Signet marigold	А	White, gold, yellow, red	May to September	Citrus; milder than T. erecta	-	Ornamental. Prefers full sun; well-drained soil but tolerates many soils.

#### (Continued)

Table 2: Recommended plants with edible flowers.

Type <sup>1</sup>	Flower color	Bloom	Taste	Fragrance	Comments and hints
Р	Yellow	May to June; fall	Bitter	-	Eat cooked only. Cool weather; full sun; tolerates wide range of soils.
Р	Pink, purple, white	July to August	Milder than leaves	-	Herb. Most creeping thymes have little flavor.
Р	Pink, lilac	June to September	Hay	Hay	Scatter florets on salad. Tolerates most soils; self sows.
А	Varies	July to August	Watercress, peppery	-	Container or in salads. Grow in full sun and well-drained soil.
A or TP	Lilac	Spring	Onion flavor	-	Best in full sun; tolerates part shade; prefers rich, moist, well-drained soil.
Р	Violet, pink, white	April to May	Sweet	Sweet	Ornamental. Candies well. Sun to part shade; moist, well-drained soil.
А	Violet, white, pink, yellow, multi-colored	May to July	Stronger than violets	_	Ornamental. Prefers cool weather; moist, rich, loamy soil but tolerates many soils.
Р	Creamy white with purple tinge	July	Hint of artichoke, slightly bitter	-	Ornamental. Full sun; well-drained, sandy soil.
	P P A A or TP P	P Yellow  P Pink, purple, white  P Pink, lilac  A Varies  A or TP Lilac  P Violet, pink, white  A Violet, white, pink, yellow, multi-colored  P Creamy white with purple	P Yellow May to June; fall P Pink, purple, white July to August P Pink, lilac June to September A Varies July to August A or TP Lilac Spring P Violet, pink, white April to May white A Violet, white, pink, yellow, multi-colored P Creamy white with purple	P Yellow May to June; fall P Pink, purple, white July to August Hay P Pink, lilac June to September A Varies July to August Pappery A or TP Lilac Spring Onion flavor  P Violet, pink, white April to May Sweet  A Violet, white, pink, yellow, multi-colored  P Creamy white with purple  May to July to Watercress, peppery Onion flavor  Sweet  May to July Stronger than violets	P Yellow May to June; fall Bitter —  Pink, purple, white July to August Milder than leaves —  Pink, lilac June to September Hay Hay  A Varies July to August Peppery —  A or TP Lilac Spring Onion flavor —  P Violet, pink, white with purple May to July Stronger than violets  P Creamy white with purple July Hint of artichoke, slightly bitter —

### References

Kendall, P. and J. Rausch. 2012. CSU Extension Fact Sheet 9.340, *Flavored Vinegars and Oils*. Available at: www.ext. colostate.edu/pubs/foodnut/09340.html. Knight A.P. *A Guide to Poisonous House* 

Knight A.P. A Guide to Poisonous House and Garden Plants. Teton New Media. 2006.

Lampe, Kenneth F. *AMA Handbook of Poisonous and Injurious Plants*, 1985. American Medical Association.